

ICEAB 2010
**Proceedings of International Conference on
Environmental Aspects of Bangladesh (ICEAB)**
September 4, 2010

<http://binbd.com/benjp/ICEAB>

Organized by

BENJapan (Bangladesh Environment Network Japan chapter)

Website: <http://binbd.com/BENJP>

E-mail: benjapan-owner@yahoogroups.com atiqahad@yahoo.com

General Chair:

Hiroyuki Miyake, University of Kitakyushu, Japan

Conference Secretary:

Md. Atiqur Rahman Ahad, University of Dhaka, Bangladesh

©ICEAB2010, BENJapan. Personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution to servers or lists, or to reuse any copyrighted component of this work in other works must be obtained from BENJapan in official writing.

Printed in Japan (*Touchiku Printing Co.* www.touchiku.co.jp)

Cover Designer: Muhammad Nurul Amin, Web: www.nurulam.in e-mail: hello@nurulam.in

ISBN: (print and electronic edition) 978-984-33-1979-1

RW04

Status of Arsenic Test and Exposure to Arsenic-contaminated Tubewell Water: A Population-based Study in Southern Bangladesh

Nepal C Dey¹, AKM Masud Rana¹, Milan K. Barua² & Babar Kabir²

¹Research and Evaluation Division, ²Disaster, Environment and Climate Change, BRAC

Abstract-This study explores the status of arsenic test of tubewell and exposures to arsenic-contaminated tubewell water in the arsenic prone southern Bangladesh. Through a survey of 6,593 households, a total of 3,812 tubewells were sampled at baseline (2006/07) and 3,591 at midline (2009). Households were selected through multi-stage sampling procedure where each sub-district was considered as a cluster. Findings reveal that proportion of unmarked (to identify presence of arsenic) tubewells increased from 80% at baseline to 90% at midline ($p<0.001$). Proportion of tested tubewells was significantly decreased from 75% at baseline to 69% at midline ($p<0.001$). Of the tested tubewells, 60% was reported to be free from arsenic contamination. Around 44% of the household members usually drink water from arsenic contaminated tubewells. This study underlines that information, education and communication campaigns should be strengthened to raise awareness regarding negative effect of arsenic on health.

INTRODUCTION

The arsenic contamination of groundwater in Bangladesh is the biggest natural calamity in the world in terms of the affected population. When rural people had developed the habit of drinking tubewell water, arsenic was found in tubewell water in many parts of Bangladesh too high concentration which has drastically reduced the coverage of safe water [1]. It was reported that the people in 62 out of 65 districts of Bangladesh were suffering from various diseases because of drinking arsenic-contaminated tubewell water. More than 70% of the country's 150 million people are at risk and 32-52 million are potentially exposed to arsenic contamination above the Bangladesh drinking water standard ($>5\mu\text{gL}^{-1}$) where about 90% people live in rural areas [2]. Around 29% of the tubewells are also contaminated with bacteria which are mainly due to poor maintenance of the tubewell surroundings. The Department of Public Health Engineering (DPHE) is responsible for marking arsenic-contaminated and arsenic-free tubewells as red and green respectively and warned the villagers against drinking water from the red tubewells.

Arsenic contamination impacts on poor population of Bangladesh. It has been well documented that the first victims of such pollution are the people with low nutrition (often people with low body weight) [3]. Women suffer from arsenic not only in terms of physical illness but also social consequences as they cannot get married and are seen as a burden to their families and their communities [4]. In a recent survey conducted in all districts of Bangladesh, nearly 19,000 arsenic patients have so far been identified

from 104.9 million arsenic-affected population. This may cause many health effects including cancer of the liver, lung, bladder and skin [5]. The United States Environmental Protection Agency (EPA) said that in Bangladesh with regard to protected future cases of arsenic-related health burden, skin cancer would affect 375,000 people [6]. The estimate also suggested that in Bangladesh approximately 6,500 people may die from cancer every year and 326,000 people in 50 years, while 2.5 million people will develop some kind of arsenicosis during that period. Availability of arsenic mitigation technologies and improving habit for using these technologies are the basic options for safe drinking in the rural areas. Recently, a report on water and sanitation explained necessity of the immediate measures against the menac of arsenic contamination in Bangladesh [7]. Acceptance of alternative arsenic mitigation technologies depends not only on the level of awareness but also physiocultural and socioeconomic variations among communities [8]. Currently 16% population is covered by arsenic mitigation technology and it is expected that the coverage will be increased to about 22% by 2015 [1].

The government of Bangladesh, incorporating stakeholders, has initiated different schemes to scale up access to safe water in rural areas and small towns. Still many rural people drink tubewell without knowing arsenic contamination level and some people drink even knowing as there is no alternate source of water available to them. Lack of access to safe water and sanitary latrine, and poor hygiene behavior are responsible for death of thousands of people of Bangladesh in every year. Water, Sanitation and Hygiene (WASH) programme of BRAC has been working with the government of Bangladesh since 2006 in 150 sub-districts for providing safe and sustainable drinking water, improving sanitation and hygiene practices to achieve Millennium Development Goals (MDG 7, target 10) by 2015 [9]. Although there are several reports published on the impact of WASH intervention. However, no study has yet been conducted on the impact of WASH intervention for safe water in the arsenic-affected areas of southern Bangladesh. Therefore, the aim of the study was to explore the changes in status of arsenic test and exposures to arsenic-contaminated tubewell water in arsenic prone southern Bangladesh.

METHODS AND MATERIALS

This study was done in 11 sub-districts under BRAC WASH programme where arsenic contamination was found

common (Table 1). Arsenic-related information was collected by using a structured questionnaire developed during the baseline followed by midline survey. Respondent was adult female member of a household who used tubewell water for drinking.

Table 1 Study sites

Name of district	Name of sub-district
Jessore	Monirampur
	Keshobpur
	Jhikorgachha
	Sharsha
Khulna	Dumira
	Fultala
	Rupsha
	Botiaghata
Noakhali	Sonaimuri
	Keshobpur

Sampling procedure for the household survey

Through survey of 6,593 households, the study sampled 3,812 tubewells in the baseline (2006/07) followed-up 3,591 in the midline (2009) in 15 sub-districts of the southern Bangladesh. Households were selected through multi-stage sampling where each sub-district was considered as a cluster. A total of 600 respondents were selected from each cluster. The economic status of the respondents was classified as ultra poor, poor and non-poor households according to WASH baseline findings [8]. The ultra poor people who were landless or homeless and had no fixed source of income were selected. The household who had up to 100 decimal of land (agricultural and homestead) and used to sell manual labour for living was considered as poor. On the otherhand, the household that do not fall in any of the above category called non-poor.

Data collection and quality control

Data were collected by the trained enumerators who had previous experience and completed at least fourteen years of education. Information on water and other demographic and socioeconomic variables for each sampling unit was collected using structured questionnaires and spot observations. After entry, 20% of the data were rechecked to identify any inconsistencies.

Statistical Analysis

The statistical analysis was performed with SPSS 16.0 software. The groups were compared for all variables using the chi-square test to compare categorical value. The differences were considered statistically significant at the $p < 0.05$ (two-tailed test) level with admissible error of 5%.

D. Ethical consideration

Before the interview verbal informed consent was taken from the participants. The verbal informed consent form was read out to the participants in native language (Bengali)

by the interviewer. Participants were informed about the objective of the study. They were also informed that their participation was entirely voluntary and they had the right to withdraw from the study at any time. Furthermore, it was informed that they had right to refuse answers to any questions if they feel uncomfortable. Confidentiality was maintained; survey questionnaire was kept secure with the researcher and was not shared with anybody other than for research purpose. Permission to conduct this study was obtained early from IRPC (Internal Review and Publication Committee) of the Research and Evaluation Division of BRAC.

RESULTS AND DISCUSSION

Profile of the respondents

The economic status, education, main occupation, marital status, responsibility to household water collection and age of the participants are presented in Table 2. Around 53% respondents were non-poor and 30% were poor and remaining were ultra poor. The main occupation of most of the respondents was household works. Around 36% respondents were illiterate and 30% passed primary education, around 32% secondary, 1.2% higher secondary and remaining passed bachelor degree. More than 90% respondents were married. Among the respondents, 43% were in age limit 10-30 years, 49% in 31-50 years and remaining were in higher ages.

Status of tubewell marking

Red-marked tubewells indicate arsenic contamination of water while green-marked tubewells indicate arsenic free. The percentages of unmarked tubewells increased significantly ($p < 0.001$) in the midline (Table 3) though it was high risk area for drinking tubewell water. This might be because of lack of awareness among the households or lack of monitoring and improper management.

Status of response on arsenic testing and results of testing tubewell water

The status of responses on arsenic testing and the results of testing in tubewell water among the study households are presented in Table 6 and 7. Proportion of tested tubewells significantly decreased from 75% at baseline to 69% at midline. However, around 24% of the tubewells were not tested at baseline and 29% at midline (Table 6). It indicates that though it was a high risk area for drinking tubewell water but people were not much aware of arsenic contamination in tubewell water and its affect on public health.

Table 2 Profile of the respondents

Indicators	Percent
Economic status	
<i>Ultra poor</i>	17.7
<i>Poor</i>	29.5
<i>Non-poor</i>	52.8
Education	
<i>Illiterate</i>	35.6
<i>Primary</i>	30.2
<i>Secondary</i>	31.7
<i>Higher secondary</i>	1.2
<i>Bachelor</i>	1.3
Main occupation	
<i>Household works</i>	93.2
<i>Day laborer</i>	1.9
<i>Student</i>	1.2
<i>Employee</i>	0.8
<i>Business</i>	0.2
<i>Others (Agriculture, rickshaw pulling, work in bus, etc.)</i>	2.7
Marital condition	
<i>Married</i>	90.8
<i>Unmarried</i>	2.1
<i>Widow</i>	5.9
<i>Separated</i>	0.9
<i>Divorced</i>	0.3
Responsible for household water collection	
<i>Women</i>	95
<i>Other members</i>	5
Age (year)	
<i>11-30</i>	43
<i>31-50</i>	49
<i>51-60</i>	6
<i>61-above</i>	2
n	6,593

Table 3 Status of tubewell marking

Status	Baseline	Midline	P value
Red marked (%)	8.9	3.8	
Green marked (%)	8.7	5.9	
Unmarked (%)	82.4	90.3	<0.001
n	3812	3591	

Table 4 shows the marking status of tubewells in different districts. Marking of tubewells was less in Khulna district both at baseline and midline where more than 90% of the tubewells were unmarked (Table 4). Whereas, most of the tubewells (96%) were unmarked in the Noakhali district, the highest arsenic concentration ($4730 \mu\text{gL}^{-1}$) area which was a new experience in 20 years in arsenic research [10].

Table 4 Status of marking of tubewells in different districts

Marking status	Arsenic contaminated tubewells					
	Jessore		Khulna		Noakhali	
	Baseline	Midline	Baseline	Midline	Baseline	Midline
Red marked	13.9	6.9	1.2	0.9	9.7	1.8
Green marked	12.8	7.7	7.5	5.9	3.0	2.2
Unmarked	73.4	85.4	91.4	93.2	87.3	95.9
n	1685	1584	1191	1246	936	761

Table 5 presents the status of marking of tubewells based on economic category of the participants. The proportion of unmarked tubewells increased in the midline among all categories (ultra poor, poor and non-poor) of households. The proportion of unmarked tubewells increased significantly from 81% at baseline to 93% at midline for ultra poor households which were found highest among all categories of households.

Table 5 Status of marking of tubewells by economic category

Marking status	Households					
	Ultra poor		Poor		Non-poor	
	Baseline	Midline	Baseline	Midline	Baseline	Midline
Red marked (%)	11.5	2.6	8.2	3.7	8.6	4.0
Green marked (%)	7.5	4.0	9.1	6.7	8.8	6.0
Unmarked (%)	81.0	93.4	82.7	89.6	82.6	89.9
n	521	454	1061	998	2230	2128

Of the tested tubewells, around 60% were reported to be arsenic-free and 40% had arsenic contamination (Table 7) which could be compared with the findings of other research where 43% of tubewells were found to be arsenic-contaminated [10]. Some changes occurred where a small proportion likely to be increased at midline who did not know about testing of tubewell water for arsenic identification and the test results. This might be because of lack of awareness raising activities and lack of facilities for arsenic testing in the study area or both. Knowledge on the affect of arsenic in public health is important for raising

awareness for safe water. It would be more effective for the people of that area if awareness raising activities could be increased on arsenic free safe water and mitigation options.

Use of arsenic contaminated tubewells water

Table 8 shows that around 44% of the households used to drink water from arsenic-contaminated tubewells in both dry and rainy seasons. This is another indication of lack of awareness on arsenic problem or lack of availability of alternative sources of safe water. The use of arsenic

contaminated water for cooking was lower (10.9%) compared to drinking.

Table 6 Status of responses on arsenic testing of tubewell water

Status of knowledge			P value
	Baseline	Midline	
Tubewells tested for arsenic identification (%)	74.9	69.3	<0.001
Tubewells not tested for arsenic identification (%)	23.5	28.6	
Don't know whether tubewells water tested/not tested (%)	1.6	2.1	
n	2212	2281	

Table 7 Status of responses on the results of testing of tubewell water for arsenic contamination

Status	Baseline	Midline	P value
Arsenic free tubewells (%)	58.0	59.8	<0.001
Arsenic contaminated tubewells (%)	41.8	39.7	
Don't know about the results of testing (%)	0.2	0.6	
n	1656	1581	

Table 8 Use of arsenic contaminated tubewell water

Purpose	Ultra poor	Poor	Non-poor	All
<i>Dry season</i>				
Drinking (%)	41.3	49.1	40.4	43.6
Cooking (%)	6.3	15.1	11.3	10.9
<i>Rainy season</i>				
Drinking (%)	41.3	49.4	40.5	43.7
Cooking (%)	8.2	19.1	14.2	13.8
	63	159	399	621

CONCLUSION

This study reveals that the household members still drink water from arsenic-contaminated tubewells. Proportion of respondents who did not know about arsenic test and test results of tubewell water increased in the midline. This might be because of lack of awareness or lack of facilities for arsenic testing in the study area where more than 95% of women were responsible for collecting water from tubewells. As proportion of unmarked tubewells was high both at baseline and midline though study area was a highly risk for drinking tubewell water, it is necessary to motivate people for marking their tubewells based on results of arsenic testing. Information, education and communication campaigns should be strengthened to raise awareness and thereby testing and marking of tubewells will be enhanced. Besides, availability of arsenic-free safe water needs to be urgently ensured to prevent health hazard.

REFERENCES

- [1] GED & UNDP, "Millennium Development Goals Needs Assessment & Coating 2009-2015 Bangladesh", General Economic Division, Planning Commission, GOB & UNDP, 2009, pp. 175-251, available: www.undp.org.bd
- [2] BRAC, "Arsenic contamination in tubewell water & its awareness among women in WASHprogramme areas: A population-based survey", unpublished.
- [3] IMF, "Bangladesh poverty reduction strategy paper", International Monetary Fund, Issues 5-410, pp. 1-183.
- [4] SOS, "Women, silent victim of groundwater poisoning", the International SOS, Accessed on February 2010 Available: www.sos-arsenic.net
- [5] A. H Smith, C Hopenhayn-Rich, M. N Bates, H. M Goeden, I. Hertz-Picciotto, H.M Duggan, "Cancer risks from arsenic in drinking water" Environ. Health Perspect, 1992, vol.97, pp. 259-267.
- [6] K. Roy, N. C. Dey and MIM. Zulfiker, "About Bangladesh and Country's Arsenic Problem: At a Glance", In: *Arsenic Calamity of Groundwater in Bangladesh: Contamination in Water, Soil and Plants*, K. Roy, Ed. Nihon University, Japan, 2008, pp. 1-11.
- [7] World Bank, "Bangladesh Country Environmental Analysis" in *Bangladesh Development Series*, 2006, Paper No: 12, Available: www.worldbank.org.bd/bds
- [8] BRAC, "The use of alternative safe water options to mitigate the arsenic problem in Bangladesh: community perspective", RED,2003, Mono. 24, pp. 1- 61.
- [9] BRAC, *WASH programme of BRAC: Towards attaining the MDG targets-baseline findings*, Research and Evaluation Division, 2008, ch.5.
- [10] D. Chakraborti, B. Das, B. Nayak, A. Pal, MM Rahman, MK Sengupta, MA Hossain, S. Ahmed, KC Biswas, M. Sahu, KC Saha, SC Mukeherjee, S. Pat, RN Dutta, and Q. Quamruzzaman, "Groundwater arsenic contamination and its adverse health effects in the Ganga-Meghna-Brahmaputra plan", In: *Arsenic Calamity of Groundwater in Bangladesh: Contamination in Water, Soil and Plants*, K. Roy, Ed. Nihon University, Japan, 2008, pp. 13-52

Author Index

A	A.T.M. Rafiqul Hoque	FB04, FB05, FM01, FM02, GE02		
	A K M Sadrul Islam	CC07		
	A. Hoshino	WM01		
	A. Salam Khan	FM04		
	A.A. Masrur Ahmed	RW08, WM04, WM05		
	A.A.M. Ahmed	RW09, WM05		
	A.B.M. Badruzzaman	GE02		
	A.F.M. Akhtaruzzaman	RW11		
	A.G. Tithi	WM05		
	A.K.M. Golam Sarwar	FB01		
	A.N.M. Fakhruddin	RW01		
	Abdullah-Al-Mahin	RW01		
	A.H.M. Kamal Sikder	PH04		
	Abu Kowsar	GE04, GE05		
	Ahammad Al Biruni	WM10		
	Ahmed, K. M. Tanvir	PA02, RW06		
	Akharuzzaman, M.	EE03		
	Akinori Jyo	RW14		
	Akio Hagihara	FM01, FM02		
	AKM Masud Rana	RW04		
Atsushi Deguchi	EE03			
B	Babar Kabir	RW04		
	Biplob K. Biswas	WM01		
C	Chamhuri Siwar	EE01, EE02		
	Chowdhury M. Alam	WM09		
D	D.A. Begum	PA02, RW06		
	Dipesh Joshi	FB06		
E	E. Ali	RW09		
F	Farhana Jabbar	FB04		
	Feroz Kabir Kazi	PA04		
	Fowzia G. Rashid Lopa	WM06		
H	H. Harada	WM01		
	H. Kawakita	WM01		
	H.M.A. Mahzuz	WM04		
	H. Rahman	FM04		
	Hajime Araki	FB01		
	Hasan M. Abdullah	WM10		
	Helena Akhter Sikder	PA03		
	I	Ishrat Islam	CC06	
J	Jeeranut Suthawaree	PA03		
K	K. Inoue	WM01		
	K. Maudood Elahi	CC01		
	K. Ohto	WM01		
	K. Omine	GE07		
	K.A. Rabbani	RW03		
	K.K. Islam	FB06		
	Kangkuso Analuddin	FM01		
	Kashif Mahmud	CC07		
	Kazi Mahmuda Tasneem	RW13		
	Kazi Rizwana Mehzabeen	GE04, GE05		
	Kenbu Teramoto	PA05		
	Khairul Hassan	RW12		
	Khalid Md. Bahauddin	GE01, PA01, PH01		
	Khan Mamun Reza	GE03		
	Kiyoshi Haraguchi	WM02		
	Kiyoshi Kurosawa	RW10		
	Koli Anar	FB07		
	Kushal Roy	CC05		
	Kyoko Sasaki	WM02		
	M	M. Ahmed	RW08	
M. Ashraf Ali		RW13		
M. Azizul Moqsud		GE07		
M. Golam Mahboob		WM10		
M. Julker Nayen		FB05		
M. Rahman		FM04		
M.A. Hannan		FM03		
M.A. Haque		PH05		
M.A. Hossain		RW03		
M.A. Yusuf		WM04		
M.A.Z. Chowdhury		RW01		
M.H. Sarkar		FM04		
M.I. Khalil		WM03		
M.J.B. Alam		RW09, WM05		
M.L. Rahman		RW03		
M.M. Rahman		RW07		
M.T.H. Chowdhury		FM03		
Magali Uono		EE04		
Md. Abdul Khaleque		RW14		
Md. Anowar Hossain Bhuiyan		EE01, EE02		
Md. Azizul Kaji	PA04			
Md. Bellal Hossain	RW05			

	Md. Hanif Uddin	PH01	Shareful Hassan	RW11
	Md. Rabiul Awwal	RW14	Sherif A. El-Safty	RW14
	Md. Sabder Ali	PA04	Shinichi Sibata	EE04
	Md. Sarwar Hossain	CC05	Shungo Kato	PA03
	Md. Shamim Uddin	RW10	Siddiqua Farah Diba	EE04
	Md. Sofikul Islam	GE04, GE05	Sk. Salahuddin	PA04
	Md. Tajuddin Sikder	CC01, CC04	Ahammad	PA04
	Md. Tawhidul I. Khan	PA05	Sobahan Mia	GE06
	Md. Zakaria Hossain	RW05	Sonya Afrin	FB03
	Milan K. Barua	RW04	Souman Rudra	GE02
	Mohammed Kamal Hossain	FB04	Subarna Sharmin	FB03
	Mohammed Mohiuddin	FB05	Sylvain R. Perret	RW02
	Mohammed Nasiruddin	PA03	T Tamotsu Kozaki	WM03
	Mukand S. Babel	RW02	Tanvir Islam	PH02
	Neelopal Adri	CC06	Tariq Md. Salah Uddin	GE01, PA01
N	Nepal C Dey	RW04	Toshinori Sakai	RW05
	Nobuyoshi Ohno	GE06	Y Yoshizumi Kajii	PA03
	Noor Danish Ahrar Mundari	WM02	Z Z. Islam	FM04
	Noriaki Seko	RW14	Zahid Hasan Mahmood	GE03, GE04, GE05
	Noriko Sato	FB06	Zahid Parvez Sukhan	EE01, FM03
	Nure Ferdousee	FB04, FB05		
O	O. Quader	FB08, FM04		
Q	Qazi Azizul Mowla	FB03		
R	R. Sauerborn	PH05		
	R.K. Mazumder	RW08		
	Rabiul Islam	EE01, EE02		
	Rahman M. Bozlur	EE04		
	Reaz A. Mullick	RW02		
S	S. Hasan	RW03		
	S. Yamamoto	PH05		
	S.B. Mahtab	RW07		
	S.K. Biswas	RW07		
	S.M. Ashekuzzaman	GE02		
	S.M. Shamimur Rahman	PH03, WM08		
	Sahadev Sharma	FM01, FM02		
	Sanzida Akhter	CC02, CC03, FB02		
	Sarder M. Yahya	CC07		
	Seichi Sato	WM03		
	Shaharuddin M. Ismail	EE01		
	Shahriar Shams	CC07, PH03, WM08		
	Shajhan Siraj	PA04		
	Shamsul Chowdhury	WM07		



BEN (Bangladesh Environment Network), Japan Chapter BENJapan

Introduction of BEN and BENJapan:

BENJapan is formed with an effort to contribute in Bangladesh's environment sectors, by the Bangladeshis living in Japan in collaboration with Bangladesh Environment Network (BEN) members from all over the world. BEN is working hand in hand with BAPA ("Bangladesh Poribesh Andolon" – Bangladesh Environment Campaign). BEN has been set up to facilitate communication among experts working on Bangladesh's environmental aspects, since July 1998.

Objectives of BEN:

- a) Gather and disseminate information about environmental issues in Bangladesh
- b) Gather and disseminate information about possible solutions to Bangladesh's environmental problems
- c) Establish and strengthen connection among various environmental organizations in Bangladesh
- d) Establish and strengthen connection between environmental organizations in Bangladesh on the one hand and international environmental organizations, on the other
- e) Formulate various policies that can and should be adopted to solve and avoid environmental problems in Bangladesh
- f) Generate public opinion in favor of policies necessary to protect environment &
- g) Persuade and assist the government to adopt and implement environment-friendly policies.

5 Principles of BEN:

1. *Self-reliance*: It refers to BEN's practice in avoiding external funds and instead of relying on members' contributions to support its activities
2. *Non-partisan character*: BEN is open to people of different political persuasions (irrespective of political views)
3. *Emphasis on RB-NRB cooperation*: Resident-Bangladeshis(RB)-Non-resident-Bangladeshis(NRB) cooperation is mutually reinforcing
4. *Non-hierarchical organization*: BEN grew up as an Internet-based organization. All BEN members have basically the same rights & opportunities in voicing their views & being heard. It has no formal office bearers
5. *Consensus decision making*: Decisions in BEN are arrived through discussion and consultation

BEN Panels on Various Environmental Problems of Bangladesh:

- Arsenic and Other Ground Water Contamination
- River and Other Water Issues
- Energy Issues
- Global Warming and Climate Change
- Economic, Legal, and Management Aspects of Environment
- Household, Industrial, and Medical Waste
- Surface Water Contamination
- Air Pollution and Other Problems of Urbanization
- Jute
- *Adibasbi* Issues
- Population Growth, Forests, and Bio-diversity

© BENJapan, ICEAB2010.

BENJapan
Bangladesh Environment Network
Japan chapter
Website: <http://binbd.com/benjp>

ISBN 978-984-33-1979-1



9 789843 319791 >