

A Comparative Analysis on Antibiotic Resistance Pattern in Environmental and Clinical Bacterial Isolates Based on Whole Cell Protein Profiling by SDS-PAGE



A DISSERTATION SUBMITTED TO BRAC UNIVERSITY IN PARTIAL
FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF
BACHELOR OF SCIENCE IN MICROBIOLOGY

Submitted by

Aneeka Nawar Fatema

Student ID: 11126003

August 2015

Microbiology Program

Department of Mathematics and Natural Sciences

BRAC University

DECLARATION

I hereby declare that the thesis project titled “**A Comparative Analysis on Antibiotic Resistance Pattern in Environmental and Clinical Bacterial Isolates Based on Whole Cell Protein Profiling by SDS-PAGE**” is my own work and the research work presented here is based on the actual and original work carried out by me. All the research work has been carried out under the supervision of Fahareen-Binta-Mosharraf, Lecturer, Microbiology Program, Department of Mathematics and Natural Sciences, BRAC University, Dhaka. Any reference to work done by any other person or institution or any material obtained from other sources have been duly cited and referenced.

(Aneeka Nawar Fatema)

Candidate

Certified

(Fahareen-Binta-Mosharraf)

Supervisor

Lecturer,

Microbiology Program,

Department of Mathematics and Natural Sciences,

BRAC University, Dhaka.

Acknowledgement

The accomplishment of my research work in pursuance of my dissertation is the very first time that I have had the exposure to this kind of experimental research work. Completion of this work needed lots of help and encouragement so as not to get frustrated at any event of failure or whenever I stumbled upon any difficulties during this elaborate time of the experimental works. I was fortunate enough to have the people around me who have provided me with all the needed support.

My regards, appreciation, indebtedness and deepest gratitude towards my respected supervisor **Fahareen-Binta-Mosharraf**, Lecturer, Microbiology Program, Department of Mathematics and Natural Sciences, BRAC University, for her efforts and patience with me throughout the period of my research work. I am extremely fortunate to have her as my supervisor for she has constantly supervised me and bestowed upon me her constructive criticism, expert guidance and has given me moral support and help whenever my steps faltered during the whole period of my research work.

I also express my gratitude towards **Prof. A. A. Ziauddin Ahmad**, Chairperson, Department of Mathematics and Natural Sciences, BRAC University, for his kind cooperation, active support and constant supervision. **Prof. Naiyyum Choudhury**, coordinator of Biotechnology and Microbiology program, Department of Mathematics and Natural Sciences, BRAC University for his excellent guidance, monitoring and constant encouragement throughout the Project. His ever so welcoming nature and the enthusiastic will to help the students has helped me not only during this research work, but throughout my entire time in BRAC University.

I would also like to thank and express my deepest gratitude to **Dr. Mahboob Hossain**, Associate Professor, Department of Mathematics and Natural Sciences, BRAC University who helped and guided me in my thesis report writing and has provided suggestions regarding the setting and execution of the experimental works, interpretation of the results and ensuing directions for the whole work with great patience. It would have been impossible to submit my thesis report without his cordial help.

Also, my appreciation goes to all my departmental teachers who have often enquired about the progress of my work and very often provided me with their valuable advice. I thank them all for this kind and affectionate gesture.

Finally, I extend my special thanks and gratitude to all my seniors and batch mates in the microbiology specialized research laboratory, who have provided good working environment by sharing their advice and encouragement to make me feel at ease even in the hardest times. They have willingly helped me out with their valuable information and boosted my efforts in finishing the task properly. Without their cordial support it would be very difficult to carry out this research work.

Aneeka Nawar Fatema

August, 2015

Dedicated

To

PROF. DR. KAZI MASIHUR RAHMAN

(Maternal Grandfather)

Abstract

This comparative study was carried out to investigate the presence of antibiotic sensitive and resistant bacteria in different environmental samples of Bangladesh with specific clinical strain obtained from ICDDR, B. The environmental samples were isolated from different locations of Dhaka city to perceive the propensity of the environmental isolates to develop antibiotic resistance. The bacterial isolates were identified on the basis of standard cultural, morphological and biochemical characteristics. Antibigram was done to identify the antibiotic susceptibility of the isolates according to Clinical and Laboratory Standards Institute (CLSI) guideline. The result of this study indicated all the environmental isolates have become resistant to the tested antibiotics, and some of them have become multi-resistant to these therapeutic agents whereas all the clinical isolates showed the opposite result. The study helps to predict the future emergence and guide the development of strategies to counteract this resistance. Therefore comprehensive survey of antibiotic resistance in the environmental bacteria is required.

This comparative study was carried out to investigate the difference in protein band of resistant strain of *E.coli* and *Staphylococcus aureus* isolated from different environmental samples of Bangladesh along with specific clinical strain obtained from ICDDR, B of *E.coli* and *Staphylococcus aureus*. The antibiotic resistance of the identified organisms was carried out by disc diffusion method with commercially available disc of Sulfamethoxazole / Trimethoprim (SXT 25), Cefoxitin (FOX 30), Pefloxacin (PEF 5), Ciprofloxacin (CIP 5), Erythromycin (E 15), Gentamicin (CN 10), Kanamycin (K 30), Streptomycin (S 10), Cefuroxime Sodium (CXM 30), Nalidixic Acid (NA 30), Oxacillin (OX 1), Chloramphenicol (C 30), Nitrofurantoin (F 300). Antibiotics have different mode of action such as cell wall synthesis inhibitors, membrane permeability alternatives, protein synthesis inhibitors and DNA synthesis inhibitors. It was further studied by using crude extraction of whole cell proteins followed by separation of proteins by means of SDS-PAGE. The environmental samples were isolated from different locations of Dhaka city to perceive the propensity of the environmental isolates to develop antibiotic resistance at different time periods of a year. The bacterial isolates were identified on the basis of standard cultural, morphological and biochemical characteristics. Antibigram was done to identify the antibiotic susceptibility of the isolates according to CLSI guideline. The result of this study indicated that there is a

specific protein band difference of the environmental isolates and the clinical isolate. The study helps to predict the future emergence and guide the development of strategies to counteract this resistance. Therefore periodic and comprehensive survey of antibiotic resistance in the environmental bacteria is required.

Chapter	Section	Title	Page
1		Introduction	1-5
	1.1	Overview	2
	1.2	Emerging pathogens	3
	1.3	Bacterial resistance to antibiotics; now and then	3
	1.4	SDS PAGE	4
	1.5	SDS	4
	1.6	Polyacrylamide Gels	4-5
	1.7	Aims and Objectives	5
2		Materials and Methods	6-23
	2.1	Working laboratory	7
	2.2	Reference Bacterial Strains	7
	2.2.1	Preparation of plating bacteria	7
	2.2.2	Confirmation of the reference strains	8
	2.2.3	Biochemical confirmation of the clinical strains	8
	2.2.4	Biochemical Identification	8-11

	2.2.5	Preparation of Stock Sample	11
	2.3	Provided Antibiotic Discs	12
	2.4	Antibiotic Susceptibility Test of the Standard Clinical Strains of Bacteria	13
	2.4.1	Preparation of McFarland Solution	13
	2.4.2	Preparation of inoculums	13
	2.4.3	Comparing with McFarland solution	14
	2.4.4	Inoculation of the MHA plates	14
	2.4.5	Placement of the Antibiotic Discs	14-15
	2.4.6	Measuring zone size	15
	2.5	Environmental sample	15-16
	2.5.1	Sample processing	16
	2.5.2	Assortment of desired colonies from nutrient agar plates	16
	2.5.3	Confirmation of the organisms by biochemical tests	17
	2.5.4	Antibiotic susceptibility test of the microorganisms from environmental samples	17
	2.6	Protein profiling	17

	2.6.1	Protein extraction	17-18
	2.6.2	Preparing the SDS PAGE Gel	18-20
	2.6.3	SDS PAGE	20-22
	2.6.4	Staining and destaining of gel	22-23
3		Results	24-54
	3.1	Conformation of clinical strains	25-30
	3.2	Confirmation of the environmental strains by biochemical tests	31-33
	3.3	Selective antimicrobial activity test by means of antibiogram method	33-45
	3.4	Comparative analysis of antibiotic susceptibility profiles of clinical and environmental strains	46-50
	3.5	Qualitative analysis of SDS PAGE result	51-54
4		Discussion	55-59
5		Conclusion	60-61
		References	62-63
		Appendix I	64-73
		Appendix II	73-76
		Appendix III	77-78

LIST OF TABLES

Serial number	Title	Page
Table 1	Respective selective media for different reference strains	8
Table 2	Provided Antibiotic Discs	12
Table 3	Source of environmental samples	16
Table 4	Cultural characteristics of clinical strains on respective selective media	25
Table 5	Standard results of biochemical tests of target strains	28
Table 6	Cultural characteristics of the environmental strains	31
Table 7	Biochemical test results of the environmental strains	32
Table 8	Antibiotic susceptibility test results for clinical strains	34
Table 9	Antibiotic susceptibility test results for <i>E. coli</i> collected from salad sample	38
Table 10	Antibiotic susceptibility test results for <i>E. coli</i> collected from bovine sample	39

Table 11	Antibiotic susceptibility test results for <i>S. aureus</i> collected from human sewage sample	40
Table 12	Antibiotic susceptibility test results for <i>S. aureus</i> collected from chicken feces	41
Table 13	Relative migration distance of protein markers and respective log of MW	52
Table 14	Relative migration distance of unknown proteins	53
Table 15	Determining molecular weight of unknown protein by interpolating the linear equation of the graph	54

LIST OF FIGURES

Serial number	Title	Page
Figure 1	Cultural characteristics of clinical strains on respective selective media	26-27
Figure 2	Biochemical tests for <i>S. aureus</i>	29
Figure 3	Biochemical tests for <i>E. coli</i>	29
Figure 4	Oxidase Test	30
Figure 5	Catalase test	30
Figure 6	Outcome of given antibiotics on clinical strain of <i>E. coli</i> K12	35
Figure 7	Susceptibility pattern of supplied antibiotics on clinical strain of <i>S. aureus</i>	36
Figure 8	Outcome of given antibiotics on <i>E. coli</i> B3 isolate collected from bovine sample	42
Figure 9	Outcome of given antibiotics on <i>E. coli</i> S3 isolate collected from salad sample	43
Figure 10	Effectiveness of applied antibiotics on <i>S. aureus</i> C3 isolate collected from chicken feces	44
Figure 11	Effectiveness of applied antibiotics on <i>S. aureus</i> H3 isolate collected from human sewage sample	45
Figure 12	Comparison among the efficiency of different antibiotics on clinical and environmental samples of <i>E. coli</i> spp collected from bovine sample	47
Figure 13	Comparison among the efficiency of different antibiotics on clinical and environmental samples of <i>E. coli</i> spp collected from salad sample	48
Figure 14	Comparison among the efficiency of different antibiotics on clinical and environmental samples of <i>Staphylococcus aureus</i> spp collected	49

	from chicken feces	
Figure 15	Comparison among the efficiency of different antibiotics on clinical and environmental samples of <i>Staphylococcus aureus</i> spp collected from human sewage sample	50
Figure 16	SDS PAGE result analysis	51
Figure 17	Graph: log MW vs. Rf curve for marker proteins	53

LIST OF ABBREVIATIONS

❖ BPA	Baird-Parker Agar
❖ CFU	Colony Forming Unit
❖ EMB	Eosin Methylene Blue Agar
❖ MAC	MacConkey Agar
❖ ICDDR,B	International Center for Diarrheal Disease Research, Bangladesh
❖ IMViC	Indole, Methyl red, Voges-Proskauer, Citrate
❖ LB	Luria Bertani
❖ MSA	Mannitol Salt Agar
❖ MR	Methyl Red
❖ MRSA	Methicillin-resistant <i>Staphylococcus aureus</i>
❖ TSB	Trypticase Soy Broth
❖ MYP	Mannitol Egg Yolk Polymyxin Agar
❖ NA	Nutrient Agar
❖ TSI	Triple Sugar Iron
❖ XLD	Xylose Lysine Deoxycholate
❖ MHA	Mueller-Hinton Agar
❖ UTIs	Urinary Tract Infections
❖ SXT 25	Sulfamethoxazole / Trimethoprim
❖ FOX 30	Cefoxitin
❖ PEF 5	Pefloxacin
❖ CIP 5	Ciprofloxacin
❖ E 15	Erythromycin
❖ CN 10	Gentamicin
❖ K 30	Kanamycin
❖ S 10	Streptomycin

❖ CXM 30	Cefuroxime Sodium
❖ NA 30	Nalidixic Acid
❖ OX 1	Oxacillin
❖ C 30	Chloramphenicol
❖ F 300	Nitrofurantoin
❖ SDS-PAGE	Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis

Chapter 1: Introduction

1.1 Overview

It has long been established that, natural selection shapes the evolution of DNA based organisms, through the course of survival or extinction within the dynamic environment of the planet. Through breakthrough scientific discoveries, the all-connecting symbiotic nature of all living things have become more and more evident. With this knowledge it is now understood far better than ever that as humans, with a huge sustained footprint on the environment, we have become a substantial influence in the context of evolution. This idea is a bit hard to grasp as most of the noticeable evolution takes place over thousands to millions of years, but when it comes to microorganisms, it is fairly observable within human lifespan that these single celled creatures are adapting to the direct or indirect interactions with us and the environment as a whole, with ferocious rapidity.

A growing interest among the community of microbiologists is to study this dynamic relationship as it is imperative to maintain our bloated life expectancy bestowed by the era of antibiotics. In this trend, common disease-causing microorganisms and our methods to eradicate or control them are subjects of repeated rigorous scientific experiments. These studies are carried out by various pharmaceutical companies, organizations for disease control, genetic engineers, biotechnologists, evolutionary microbiologists and every other group that have a vested interest in improving and maintain the quality of human life.

A stark example can be the evolution of the enteric disease causing pathogens throughout the history of modern medicine. In the public consciousness, a disease like diarrhea has gone from being a sweeping epidemic that leaves us mostly helpless, to a benign inconvenience that we have to endure for a few days, but this underlies an uphill battle between synthetic antibiotics and disease causing pathogens, that microbiologists have been fighting ever since the days of penicillin.

1.2 Emerging pathogens

A humongous number of organisms are associated with a variety of infectious diseases, and are considered to be human pathogens. Not only are the pre existing pathogens menacing the human health, but also some pathogens have only been recognized very recently. Such pathogens are largely accountable for a massive number of deaths every year. As many new diseases are held responsible for this global burden, which is taking its toll on human health, it is also true for diseases that had been declining over the past few years [1]. The re emergence of these diseases is associated with certain transmission routes and also different and wide range of hosts.

1.3 Bacterial resistance to antibiotics; now and then

Antibiotics, since its invention have been vital in the fight against infectious diseases that are caused by various bacteria as well as other microbes. A matter of great misery is that although we have all these advanced antibiotics now at our disposal, reports have shown that no antibiotic can actually last effective for too long. As a result, resistance in the pathogens to common therapeutic agents is increasing in recent years. Thus, a mere cultural contaminant or a harmless microbe can eventually become great human pathogen over time [2].

The abundance and indiscriminate use of the commercial antimicrobial drugs that are commonly used in the treatment of infectious diseases is the leading reason of multiple drug resistance in human pathogenic microorganisms that has been developed over the past years (Table 1) [5]. This development of antibiotic resistance has a number of favoring factors, those including the characteristics of the hosts, the usage of antibacterial agents, the specific nature of the relationship of bacteria to antibiotics, as well as environmental factors [3]. Since the bacterial pathogens have the ability to rapidly evolve, this helps them to come up with new ways to dodge the host defenses and become resistant to the antibiotic treatments. It is a matter of great concern that more and more pathogens are showing resistance to multiple drugs. In addition the high levels of antibiotics used in humans and animals now a day have amplified the emergence of these antibiotic resistant strains that are the main concern of this study [4].

1.4 SDS-PAGE

The full form of SDS-PAGE is Sodium Dodecyl Sulfate the purpose of SDS-PAGE is to separate proteins according to their size, and no other physical feature. The name itself will explain the entire reasoning of SDS-PAGE, the point is to understand in details the two names SDS and PAGE accordingly.

1.5 SDS

The main goal is to separate different protein molecules of different shapes and sizes, the first thing to do is to denature the proteins. So that there is no longer have any secondary, tertiary or quaternary structure to have a linear form.

The separation of macromolecules is done in an electric field is called electrophoresis. A very common method for separating proteins by electrophoresis uses a discontinuous polyacrylamide gel as a support medium and Sodium Dodecyl Sulfate (SDS) to denature the proteins..

It is an anionic detergent, meaning that when dissolved its molecules have a net negative charge within a wide pH range. A polypeptide chain binds amounts of SDS in proportion to its relative molecular mass. The negative charges on SDS destroy most of the complex structure of proteins, and are strongly attracted toward an anode (positively-charged electrode) in an electric field.

1.6 Polyacrylamide gels

Polyacrylamide gels restrain larger molecules from migrating as fast as smaller molecules. Because the charge-to-mass ratio is nearly the same among SDS-denatured polypeptides, the final separation of proteins is dependent almost entirely on the differences in relative molecular mass of polypeptides. In a gel of uniform density the relative migration distance of a protein (R_f , the f as a subscript) is negatively proportional to the log of its mass. If proteins of known mass are run simultaneously with the unknowns, the relationship between R_f and mass can be plotted, and the masses of unknown proteins estimated.

Protein separation by SDS-PAGE can be used to estimate relative molecular mass, to determine the relative abundance of major proteins in a sample, and to determine the distribution of

proteins among fractions. The purity of protein samples can be assessed and the progress of a fractionation or purification procedure can be followed. Different staining methods can be used.

1.7 Aims and objectives

Resistance to antimicrobial agents has resulted in morbidity and mortality from treatment failures and increased health care costs. Even though defining the precise public health risk and estimating the increase in costs is not a simple undertaking, there is little doubt that emergent antibiotic resistance is a serious global problem. Widespread antibiotic usage exerts a selective pressure that acts as a driving force in the development of antibiotic resistance.

Although pharmaceutical industries have produced a number of new antibiotics in the last few decades but simultaneously the resistance of the microorganisms to these drugs have also increased. Due to such increasing resistance in microbes and synthetic antibiotic side effects, it is now necessary to access resistance pattern of pathogenic organisms against standard antibiotics to develop therapeutic alternatives.

On the basis of above context, the objectives of the present study are:

1. The degree of antibacterial activity will be evaluated considering the diameter (mm) obtained for the zone of inhibition on each of the replicate agar plate.
2. The resistance pattern of both clinical and environmental samples will be recorded for a comparative analysis.
3. The purpose of SDS-PAGE is to separate proteins according to their size, and no other physical feature.

Chapter 2: Materials and Methods

2.1 Working laboratory

All project works were performed in the Microbiology and Biotechnology Laboratory, Department of Mathematics & Natural Sciences, BRAC University.

2.2 Reference Bacterial Strains

In this study, five standard clinical strain of *E. coli K12* and *Staphylococcus aureus*, were used. All these species were obtained from ICDDR, B (International Center for Diarrheal Disease Research, Bangladesh).

2.2.1 Preparation of plating bacteria

- A single bacterial colony of each clinical isolates was inoculated into a screw cap test tube containing 5 ml Luria Bertani (LB) broth.
- The tube was incubated at 37°C in a shaking incubator (Daihan Scientific, Korea) at 120 rotations per min (rpm) for 3 hours.
- After incubation 1 ml culture from the test tube was taken in an eppendorf and centrifuged (Eppendorf, Germany) at 12,000 rpm for 10 min and the supernatant was discarded.
- Bacterial pellet was suspended in 1 ml of 10 mM MgSO₄ properly to make it homogenous.
- The suspension was stored at 4°C until used.
- The suspensions were then diluted 1:100 in TSB broth to obtain 10⁶ CFU/ml before use.

2.2.2 Confirmation of the reference strains

Reference bacterial strains were identified routinely to distinguish each organism by sub culturing on recommended selective media (Table 1). The cultural properties of each organism were observed and recorded.

Table 1: Respective Selective Media for Different Reference Strains

Bacterial Strain	Selective Media
<i>E .coli</i> K12	MacConkey agar and Eosin-Methylene blue (EMB) agar
<i>Staphylococcus aureus</i>	Mannitol Salt agar (MSA)

2.2.3 Biochemical confirmation of the clinical strains

Subsequently each bacterial clinical strain were employed for morphological and biochemical confirmation tests. Recommended biochemical tests like Methyl red test, Voges–Proskauer test, Indole test, Citrate utilization test, Oxidase test, Catalase test, TSI agar test were performed for each of the strains.

2.2.4 Biochemical Identification

Biochemical tests were performed with specific standard isolates developed in specific media according to the standard methods described in Microbiology Laboratory Manual [6]. Before starting the process of any biochemical identification test, all the bacterial cultures were grown on nutrient agar plates in the incubator at 37°C.

Methyl red (MR) test

- Bacterium to be tested was inoculated into 3 ml dextrose phosphate broth (MR-VP broth), which contained dextrose and a phosphate buffer and incubated at 37°C for 24 hours.
- Over the 24 hours, the mixed-acid producing organism might produce sufficient acid to overcome the phosphate buffer and remain acidic.
- The pH of the medium was tested by the addition of five drops of MR reagent. Development of a red color was taken as a positive result. MR negative organism would give a yellow color.

Voges–Proskauer test

- Bacterium to be tested was inoculated into 3 ml dextrose phosphate broth (MR-VP broth) and incubated at 37°C for 24 hours.
- To the aliquots of each broth cultures 10 drops of Barritt's reagent A was added and the cultures were shaken.
- Immediately, 10 drops of Barritt's reagent B was added and the cultures were shaken again.
- Cultures were then kept aside for 15 minutes for the reaction to occur.
- After 15 minutes, the colors of the cultures were examined and the results were recorded. Appearance of a red color was taken as a positive result.

Indole test

- Bacterium to be tested was inoculated in 6 ml peptone water, which contains the amino acid tryptophan and incubated at 37°C for 24 hours.
- Following overnight incubation, five drops of Kovac's reagent were added.
- Then the colors of the cultures were examined and the results were recorded. Formation of a rose red ring at the top indicates a positive result. A negative result can have a yellow or brown layer.

Citrate utilization test

- A single bacterial colony of each bacterium to be tested was picked up from each nutrient agar plates by a needle and inoculated into the slope of Simmon's citrate agar and incubated at 37°C for 24 hours.
- Over these 24 hours, the organism which had the ability to utilize citrate would change the color of the media.
- All the media that changed their color from green to a Prussian blue, is taken as a positive result. A negative slant would have no growth of bacteria and would remain green.

Triple Sugar Iron (TSI) Test

- A single bacterial colony of each bacterium to be tested was picked up from each nutrient agar plates by a needle and stabbed into the TSI containing dextrose, lactose and sucrose butt.
- Caps of the tubes were loosened and incubated at 35°C for overnight and were examined after 18-24 hours for carbohydrate fermentation, CO₂ and H₂S production.
- A yellow (acidic) color in the butt indicated that the organism being tested is capable of fermenting all the three sugars, whereas a red (alkaline) color in the slant and butt indicated that the organism being tested is a non fermenter.
- A black precipitation in the butt of the tube is the indication of H₂S production.
- Presence of bubbles, splitting and cracking of the medium is the indication of CO₂ gas production.

Oxidase test

- Two drops of oxidase reagent (*p*-Aminodimethylaniline oxalate) were added onto the filter paper (Whatman, 1MM).
- A loopful of each bacterium to be tested were taken from nutrient agar plate and streaked onto the filter paper (Whatman, 1MM).
- A positive reaction would turn the paper from violet to purple within 1 to 30 seconds. Delayed reactions should be ignored as that might give false positive result.

Catalase test

- One drop of catalase test reagent (hydrogen peroxide) was placed on a sterile glass slide.
- An isolate from a nutrient agar plate was picked up with a sterile toothpick and placed on to the reagent drop.
- This was done with each of the bacterium to be tested.
- An immediate bubble formation indicates a positive result.

2.2.5 Preparation of Stock Sample

Short term preservation

3 ml of T₁N₁ agar butt in each vial was inoculated by stabbing isolates from nutrient agar plates. Then the vials were incubated at 37°C (SAARC) for 6 hours. After incubation, the surface of the medium was covered with 200 µl sterile paraffin oil and the vials were stored at room temperature appropriately labeled

Long term preservation

For long-term preservation, 500 µl of bacterial culture grown in Trypticase Soy Broth at 37°C for 6 hours was taken in a sterile cryovial. Then 500 µl of sterile glycerol was added to the broth culture and the cryovial was stored at -20°C.

2.3 Provided Antibiotic Discs

In this study, the effectiveness of thirteen different antibiotics was determined. They are listed below in the table:

Table 2: Provided Antibiotic Discs

Antibiotic	Disc Identification Number
Sulfamethoxazole / Trimethoprim	(SXT 25)
Cefoxitin	(FOX 30)
Pefloxacin	(PEF 5)
Ciprofloxacin	(CIP 5)
Erythromycin	(E 15)
Gentamicin	(CN 10)
Kanamycin	(K 30)
Streptomycin	(S 10)
Cefuroxime Sodium	(CXM 30)
Nalidixic Acid	(NA 30)
Oxacillin	(OX 1)
Chloramphenicol	(C 30)
Nitrofurantoin	(F 300)

2.4 Antibiotic Susceptibility Test of the Standard Clinical Strains of Bacteria

The standard clinical strains of bacteria were tested for their sensitivity against thirteen standard antibiotics, as mentioned previously in Table 2.

2.4.1 Preparation of McFarland Solution

McFarland solution is an essential material needed before testing the microorganisms for their sensitivity. McFarland standards are used as reference to adjust the turbidity of any given bacterial suspension. This is done to make sure that the number of bacteria is within a given range to standardize the microbial testing. This would also help avoid any error in result, because if the suspension is too heavy or too diluted, an erroneous result might occur for any given anti microbial agent, which in this study, is antibiotic discs.

- 97% H₂SO₄ was diluted to a concentration of 1%
- BaCl₂.2H₂O was diluted to a concentration of 1.175%
- To make 5 ml McFarland solution, 4975 µl H₂SO₄ was mixed with 25µl BaCl₂.2H₂O.

2.4.2 Preparation of inoculums

- Using a sterile inoculating loop, one or two colonies of the organism to be tested were taken from the subculture plate.
- The organism was suspended in 6 ml of physiological saline.
- The test tube containing the saline was then vortexed to create an overall smooth suspension.

2.4.3 Comparing with McFarland solution

- Using the Colorimeter (Labtronics; ISO 9001: 2008 Certified), the OD of the McFarland solution was measured to be 11.
- Then the OD of each of the inoculums that were made was also measured with the Colorimeter.
- Only the solutions with the OD that matched with that of the McFarland solution were taken.
- The solutions that gave an OD higher than the standard solution were diluted with solution to match the standard.
- Once all the OD of the inoculums was matched with the standard, they were ready to be inoculated on MHA (Muller-Hinton Agar) plates.

2.4.4 Inoculation of the MHA plates

- A sterile swab was dipped into the inoculum tube. The swab was rotated against the side of the tube above the fluid level, using firm pressure, to remove excess fluid, but the swab was not dripped wet.
- The dried surface of a MHA plate was inoculated by streaking the swab four to six times over the entire agar surface; the plate was rotated approximately 60 degrees each time to ensure an even distribution of the inoculums.
- The plate was rimmed with the swab to pick up any excess liquid.
- Leaving the lid slightly ajar, the plate was allowed to sit at room temperature at least 3 to 5 minutes for the surface of the agar plate to dry before proceeding to the next step.

2.4.5 Placement of the Antibiotic Discs

- The forceps was sterilized by immersing the forceps in alcohol then igniting.
- Four sterile discs were placed on the surface of an agar plate, using that forceps.

- The discs were gently pressed with the forceps to ensure complete contact with the agar surface.
- Placing discs close to the edge of the plates was avoided as the zones will not be fully round and that can be difficult to measure.
- Once all discs were in place, the plates were inverted, and placed in a 37°C incubator for 24 hours.

2.4.6 Measuring zone size

- Following the incubation, the zone sizes were measured to the nearest millimeter using a ruler.
- All measurements were made with the unaided eye while viewing the back of the Petri dish.
- The zone size was recorded on the recording sheet.

2.5 Environmental sample

Different types of environmental samples were collected from different sources in the month of October, 2014 (Table 3). Throughout the month of October, the daytime temperatures had generally reached highs of around 30°C, which is about 86°F. At night, the average minimum temperature dropped down to around 21°C, which is 70°F. The average daily relative humidity for October was around 84%.

Table 3: Source of environmental samples

Environmental Sample	Collected from
Chicken feces	a chicken farm located at Banani
Cow dung	a farm located at Gulshan
Handmade salad	a restaurant located at Mohakhali
Human sewage	The river Buriganga

2.5.1. Sample processing

Six fold serial dilutions were done with the different environmental samples following the standard sample processing technique.

- 1 ml of sample suspension was taken and mixed with 9 ml of physiological saline in a test tube and the tube was vortexed properly to make a smooth and even suspension.
- Then the sample was diluted six times to get the desired concentration.
- The same procedures were followed for all four of the samples.

2.5.2. Assortment of desired colonies from nutrient agar plates

The maximum diluted sample was spread on nutrient agar plate for total count. Multiple colonies from each of the nutrient agar plates were selected depending on the cultural characteristics and transferred to specific standard selective media for further confirmation.

2.5.3 Confirmation of the organisms by biochemical tests

Multiple selected presumptive colonies from the specific media were confirmed by biochemical tests following standard protocol [6], as was previously done with the reference strains.

2.5.4 Antibiotic susceptibility test of the microorganisms from environmental samples

All the steps and methods that were followed during the antibiotic susceptibility test of the reference bacterial strains were repeated with the microorganisms now isolated from the environmental samples.

2.6 Protein profiling

The environmental strain from the sample strains that were found to be resistant to the specific antibiotics were then further sub-cultured in nutrient agar medium before the protein extraction was undertaken. The culture plates were incubated overnight at 37 ° C.

2.6.1 Protein Extraction:

Luria Bertani broth of 5ml was prepared for each strain.

2.6.1 Protein Extraction:

- Luria Bertani broth of 5ml was prepared for each strain.
- Desired sample was then inoculated into each tube from fresh subculture.
- Incubation period was maintained for 37 ° C for 24 hours.
- After incubation, the culture was taken into micro centrifuge tube of 1ml after vortexing.
- Then it was micro centrifuged at 14,000 rpm for 5 minute.

- Suspended pellet was found in the ground of the tube,
- The supernatant was discarded.
- 4x SDS dye of 20 μ l was added in each tube.
- The butt of the tube was tapped vigorously to dissolve the pellet and boiled at 100 ° C for 3 minutes.
- Lastly, the protein was preserved at -20 ° C.

2.6.2 Preparing SDS-PAGE gels:

Step 1:

- The plates and combs must be clean
- For each gel, one short plate is needed
- One spacer plate and one comb is also required
- A little bit of 70% ethanol is sprayed on the plates and wipe dry with Kimwipes.
- The combs thoroughly washed with tap water.
- It is critical to remove all dust and small particles, especially any bits of left-over polyacrylamide.

Step 2:

- The plates are set on the rack.
- The short plate on the spacer plate is layered with the spacers in between and
- The two plates are slid into the holder.
- The bottom edges of the two plates are flushed to avoid leakage.
- The plates are locked in and the holder is placed on the rack, with the bottom edges of the plates pushed into the foam/ rubber pad to make a water-tight seal.
- Test the seal is tested by pipetting or squirting a small volume of water between the plates and making sure there is no leakage.
- Blot is dried with filter paper.

Step 3:

- The separating gel is poured
- The table below is used as a guide to calculate the total volumes needed
- Pipette solutions are placed in order
- Bubbles must be avoided, which as a result will inhibit polymerization.
- The solution is then gently swirled to mix thoroughly after addition of each component.

	7.5% gel	10% gel	12.5% gel	15% gel	18% gel
ddH ₂ O	2.81 mL	2.50 mL	2.19 mL	1.88 mL	1.50 mL
40% acrylamide/bis stock	0.94 mL	1.25 mL	1.56 mL	1.88 mL	2.25 mL
1.5 M Tris, pH 8.8	1.25 mL	1.25 mL	1.25 mL	1.25 mL	1.25 mL
10% ammonium persulfate	50 µL	50 µL	50 µL	50 µL	50 µL
TEMED	5 µL	5 µL	5 µL	5 µL	5 µL
TOTAL VOLUME	~5 mL	~5 mL	~5 mL	~5 mL	~5 mL

- Once TEMED is added, the gel will begin to polymerize
- The work has to be done fast but carefully
- The gel is pipette and mixed between the plates
- There has to be enough space at the top for the stacking gel and comb.
- Layer water on top of the gel solution is carefully poured

- Once the gel has polymerized (about 10-15 mins), the top of the gel is washed off with water
- With a filter paper excess water is carefully blotted
- The gel must not be disturbed

Step 4:

- The stacking gel is poured.
- For each gel, the solutions carefully pipetted and swirled to mix after addition of each component
- The gel mix by Pipetting between the plates up to just below the edge of the short plate.
- The comb is carefully placed
- Once the gel has polymerized, slowly the comb is removed under running water
- The wells are carefully washed to avoid distorting them

ddH ₂ O	3.13 mL
40% acrylamide/bis stock	0.62 mL
1.5 M Tris, pH 6.8	1.25 mL
10% ammonium persulfate	50 µL
TEMED	5 µL
TOTAL VOLUME	~5 mL

Step 5:

- The gel is storage
- The comb is returned to storage and the gel is sandwiched between two wet pieces of paper towels.
- The gel can be stored horizontally at 4 °C for up to 1 week.

2.6.3 SDS-PAGE

Pouring the resolving gel

- The glass plates were cleaned with soap and water, then with ethanol.

- The glass plates and spacers were assembled
- A pasteur pipette was used and the bottom of the plates were sealed by spreading 1% agarose in SDS PAGE electrophoresis buffer along the bottom of the glass plates so it moves up the crack by capillary action.
- The degassed resolving gel is transferred to a beaker and APS/TEMED is added.
- It was then mixed
- The resolving gel solution was quickly added to the center of the plates to a height about 4 cm from the top for the large plates.
- Isobutanol/ absolute ethanol was quickly added to the top of this until the level reaches the top of the plates.
- Isobutanol/ absolute ethanol will prevent oxygen from getting into the gel which would oxidize it and inhibit polymerization.
- The resolving gel would polymerize in 30 minutes.
- The stacking gel reagents are added to a flask (without addition of APS & TEMED) and degas for ten minutes.

- **Pouring the stacking gel**
- Isobutanol/ absolute ethanol was poured off.
- The watery interpolate surface was dried with a piece of Whatmann paper.
- The stacking gel was polymerized,
- APS/TEMED was added,
- It was then mixed
- The polymerized resolving gel was poured on top

- The comb was inserted straight on down,
- Then a little more stacking gel was poured on the sides of the comb to fully seal it
- Any bubbles were removed from underneath the comb
- If possible, the comb was gently moved from side to side so the bubbles get into the space in between and float up
- The stacking gel polymerized in 20 to 30 minutes

Load the gel

- Tris-glycine electrophoresis buffer was poured into the upper and lower chambers
- Bubbles trapped were removed at the bottom of the glass plates in the large gel with a syringe.
- The wells were flushed with a syringe just before loading to get rid of any unpolymerized polyacrylamide that may seep in
- When loading the gel, sample was loaded in every lane and the dye front migrated more evenly
- Large thin gels were ran at a constant current of 12-15 mA.
- After the dye front entered the resolving gel, the current was turned up to 30 mA.
- The gel was stopped depending on how big the smallest protein was that was desired to visualize
- If waited for the dye front to just flows out of the gel, it would take about 2.5 - 3 hours for a large, thin gel to run.
- A thick gel would take 6 - 7 hours

2.6.4 Staining and destaining of gel

Reagents needed:

Coomassie Stain solution

Destain solution

Directions:

- The SDS-PAGE gel was removed from glass and rinsed once in ddH₂O in a suitable container with a lid.
- Coomassie Stain was added to cover the gel by 1/2 inch (~ 1.5 cm).
- The gel was incubated in the Coomassie stain for 35 minutes at 45°C in water bath.
- The Coomassie Stain was poured off. The Coomassie Stain can be recycled a couple of

times by filtering it.

- The gel was rinsed twice in ddH₂O to remove Coomassie Stain from the container.
- Fresh Distaining solution was added to cover the gel by 3/4 inch (~ 2 cm).
- The gel was incubated in the Distaining solution for 1 hour in a water bath.
- The gel was kept in the distaining solution overnight at room temperature.
- The used Destain solution can be recycled a couple of times by storing it in a sealed container

Chapter 3: Results

3.1 Conformation of clinical strains

Clinical strain of the five bacterial species i.e. *Escherichia coli* strain K12 and *Staphylococcus aureus* obtained from ICDDR,B (International Center for Diarrheal Disease Research, Bangladesh) were preliminary identified by their cultural properties (Table 4) upon streaking in the respective selective media (Table 1). Selective medium types are formulated to support the growth of one group of organisms, but inhibit the growth of another. These media contain antimicrobials, dyes, or alcohol to inhibit the growth of the organisms that are not targeted for study.

Table 4: Cultural characteristics of clinical strains on respective selective media

solates/ Organism	Cultural Characteristics						
	Medium	Size	Margin	Elevation	Form	Pigment	Consistency
<i>S. aureus</i>	MSA	Moderate (2-3mm)	Entire	Convex	Circular	Yellow	Smooth
<i>E. coli</i> K12	MAC	Large (2-3mm)	Entire	Raised	Circular	Pink	Smooth
	EMB	Large (2-3mm)	Entire	Slightly raised	Circular	Blue-black colonies with metallic green sheen	Shiny, smooth



Figure1a) *Escherichia coli*
K12 in MacConkey agar




Figure1b) *Escherichia coli*
K12 in Eosin Methylene blue
agar



Figure 1c) *Staphylococcus aureus*
in Mannitol Salt agar

Figure 1: Cultural characteristics of clinical strains on respective selective media:

- a) *Escherichia coli* K12 in MacConkey agar,
- b) *Escherichia coli* K12 in Eosin Methylene blue agar,
- c) *Staphylococcus aureus* in Mannitol Salt agar,

Isolates shown in Figure 1 showed patterns of biochemical reactions that are typical for each strain of bacteria according to Microbiology Laboratory Manual [6]. In Table 5 below, the results of biochemical tests of the clinical isolates are mentioned.

Table 5: Standard results of biochemical tests of target strains

Isolate/Organism	Biochemical Tests									
	Indole production test	Methyl red reaction test	Voges Proskauer reaction test	Citrate utilization test	TSI fermentation				Catalase activity test	Oxidase activity test
					Slant	Butt	CO ₂	H ₂ S		
<i>Staphylococcus aureus</i>	-	+	-	-	A	A	+	-	+	-
<i>E. coli</i> K12	+	-	-	+	K	K	+	-	-	-

KEY: A= acidic condition, K= alkaline condition, + = positive, - = negative, AG= both acid & gas production.




Figure: 2 Biochemical tests for *Staphylococcus aureus*




Figure: 3 Biochemical tests for *E. coli*




Figure: 4 Oxidase Test




Figure: 5 Catalase Test

3.2 Confirmation of the environmental strains by biochemical tests

Desired presumptive environmental strains of *Escherichia coli* strain K12 and *Staphylococcus aureus* are selected from specific media (Table 1) depending on their cultural characteristics (Table 6).

Table 6: Cultural characteristics of the environmental strains

Isolates/ Organisms	Cultural Characteristics						
	Medium	Size	Margin	Elevation	Form	Pigment	Consistency
Sta 1	MSA	Moderate (2-3mm)	Entire	Raised	Circular	Yellowish	Smooth
Sta 2	MSA	Moderate (2-3mm)	Entire	Convex	Circular	Yellow	Smooth
Sta 3	MSA	Large (2-3mm)	Entire	Slightly raised	Irregular	Colorless	Creamy
Sta 4	MSA	Small (1-2mm)	Entire	Convex	Circular	Colorless	Smooth
Esc 1	MAC	Small (1-2mm)	Entire	Flat	Circular	Pink	Rough
	EMB	Moderate (2-3mm)	Entire	Flat	Circular	Pink	Smooth
Esc 2	MAC	Small (1-2mm)	Entire	Raised	Circular	Purple	Smooth
	EMB	Small (1-2mm)	Entire	Raised	Circular	Pink	Smooth
Esc 3	MAC	Small (1-2mm)	Entire	Slightly raised	Circular	Slight Pink	Rough
	EMB	Moderate (2-3mm)	Entire	Flat	Circular	Colorless	Smooth
Esc 4	MAC	Large (2-3mm)	Entire	Raised	Circular	Pink	Smooth
	EMB	Large (2-3mm)	Entire	Slightly raised	Circular	Blue-black colonies with metallic green sheen	Shiny, smooth

A total of twenty colonies belonging to two target strains isolated from different environmental sources (Table 3) were selected depending on the presumptive cultural characteristics which were further confirmed by standard biochemical tests showed below in Table 7.

Table 7: Biochemical test results of the environmental strains

Isolate/Organism	Biochemical Tests									
	Indole production test	Methyl red reaction test	Voges Proskauer reaction test	Citrate utilization test	TSI fermentation				Catalase activity test	Oxidase activity test
					Slant	Butt	CO ₂	H ₂ S		
Sta 1	+	-	+	-	A	K	+	+	+	+
Sta 2	-	+	-	-	A	A	+	-	+	-
Sta 3	-	-	+	-	A	K	+	-	+	-
Sta 4	-	-	+	-	K	K	-	-	+	-
Esc 1	+	-	-	+	K	K	+	-	-	-
Esc 2	-	+	-	-	K	A	-	-	+	-
Esc 3	-	-	-	+	K	A	+	-	+	-
Esc 4	+	-	-	+	K	K	+	-	-	-

According to Table 6 and 7 Sta 2 and Esc 4 showed standard cultural characteristics as well as standard biochemical test results following Table 4 and 5, hence they were selected for further studies.

3.3 Selective antimicrobial activity test by means of antibiogram method

The standard disc diffusion test was done with all the provided antibiotics (Table 3) against five clinical and isolates of five selected environmental strains to identify their resistance pattern. The interpretive categories were defined according to the zone diameter of inhibition.

All the clinical strains showed significant susceptibility to all the antibiotics except for OX1 (Table 8). *E. coli* K12 showed highest levels of susceptibility to CIP5 which was measured to be 40mm. In case of *Staphylococcus aureus*, the maximal level of vulnerability was to PEF5, which was around 32mm in diameter. Antibiotic susceptibility test results for clinical strains are represented in Table 8 and Figures 6, 7, 8, 9, and 10 below.

Table 8: Antibiotic susceptibility test results for clinical strains

Antibiotics	Clinical Strain	
	<i>E. coli</i> K12	<i>S. aureus</i>
SXT 25	30 mm	28 mm
FOX 30	28 mm	30 mm
PEF 5	32 mm	32 mm
CIP 5	40 mm	28 mm
E 15	9 mm	9 mm
CN 10	20 mm	20 mm
K 30	19 mm	20 mm
S 10	14 mm	14 mm
CXM 30	25 mm	26 mm
NA 30	24 mm	25 mm
OX 1	0 mm	0 mm
C 30	24 mm	26 mm
F 300	26 mm	24 mm




Figure: 6a




Figure: 6b

Figure 6: (a) and (b) Outcome of given antibiotics on clinical strain of *E. coli* K12




Figure: 7a




Figure: 7b

Figure 7: (a) and (b) Susceptibility pattern of supplied antibiotics on clinical strain of *S. aureus*

A significant level of resistance was observed when all the standard antibiotics were applied against several isolates of selected environmental strains that *E. coli* was collected from salad sample and bovine sample (cow dung), and *Staphylococcus aureus* was collected from human sewage sample and chicken feces. They all showed resistance to at least one or more than one antibiotics.

The clinical strain of *E. coli* was significantly vulnerable to all the antibiotics, yet three different isolates of *E. coli* collected from salad sample showed resistance to three different antibiotics each (Table 9) and four isolates collected from the bovine sample (cow dung) showed resistance to three and four different antibiotics each (Table 10).

In case of environmental strain of *Staphylococcus aureus*, the result was astounding. Its clinical strain was susceptible to all the test antibiotics but one, having large zones of inhibition measured to be as high as 28-32mm in diameter and yet different isolates of *S. aureus* collected from human sewage sample (Table 11) and chicken feces (Table 12) showed resistance to as many as four and eight different antibiotics as shown below.

Table 9: Antibiotic susceptibility test results for *E. coli* collected from salad sample

Antibiotics	<i>E. coli</i> From Salad Sample								
	<i>E. coli</i> S1	<i>E. coli</i> S2	<i>E. coli</i> S3	<i>E. coli</i> S4	<i>E. coli</i> S5	<i>E. coli</i> S6	<i>E. coli</i> S7	<i>E. coli</i> S8	<i>E. coli</i> S9
SXT 25	8 mm	12 mm	Resistant	14 mm	14 mm	10 mm	12 mm	19 mm	12 mm
FOX 30	6 mm	19 mm	8 mm	16 mm	19 mm	12 mm	17 mm	21 mm	13 mm
PEF 5	Resistant	13 mm	12 mm	15 mm	22 mm	12 mm	14 mm	12 mm	28 mm
CIP 5	20 mm	24 mm	28 mm	24 mm	18 mm	21 mm	19 mm	20 mm	22 mm
E15	18 mm	14 mm	9 mm	14 mm	21 mm	10 mm	11 mm	13 mm	Resistant
CN 10	Resistant	10 mm	8 mm	14 mm	19 mm	17 mm	13 mm	18 mm	17 mm
K 30	14 mm	18 mm	Resistant	22 mm	22 mm	27 mm	24 mm	22 mm	Resistant
S 10	22 mm	24 mm	13 mm	19 mm	15 mm	28 mm	24 mm	20 mm	21 mm
CXM 30	Resistant	18 mm	Resistant	17 mm	11 mm	14 mm	28 mm	19 mm	15 mm
NA 30	10 mm	13 mm	6 mm	11 mm	9 mm	8 mm	18 mm	15 mm	9 mm
OX 1	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm
C 30	13 mm	9 mm	10 mm	13 mm	15 mm	18 mm	19 mm	8 mm	14 mm
F 300	16 mm	14 mm	13 mm	19 mm	11 mm	15 mm	13 mm	11 mm	12 mm

Table 10: Antibiotic susceptibility test results for *E. coli* collected from bovine sample

Antibiotics	<i>E. coli</i> From Bovine Sample								
	<i>E. coli</i> B1	<i>E. coli</i> B2	<i>E. coli</i> B3	<i>E. coli</i> B4	<i>E. coli</i> B5	<i>E. coli</i> B6	<i>E. coli</i> B7	<i>E. coli</i> B8	<i>E. coli</i> B9
SXT 25	7 mm	12 mm	Resistant	16 mm	Resistant	18 mm	12 mm	10 mm	14 mm
FOX 30	5 mm	9 mm	12 mm	16 mm	8 mm	11 mm	16 mm	14 mm	17 mm
PEF 5	Resistant	15 mm	Resistant	17 mm	12 mm	10mm	19 mm	14 mm	12 mm
CIP 5	16 mm	13 mm	8 mm	10 mm	28 mm	23 mm	10 mm	8 mm	14 mm
E15	Resistant	19 mm	9 mm	11 mm	9 mm	10 mm	17 mm	Resistant	12 mm
CN 10	12 mm	12 mm	15 mm	11 mm	8 mm	15 mm	18 mm	Resistant	14 mm
K 30	20 mm	18 mm	18 mm	10 mm	Resistant	22 mm	11 mm	16 mm	19 mm
S 10	Resistant	18 mm	13 mm	17 mm	13 mm	24 mm	11 mm	20 mm	16 mm
CXM 30	0 mm	13 mm	16 mm	16 mm	20 mm	22 mm	19 mm	Resistant	14 mm
NA 30	0 mm	0 mm	Resistant	0 mm	6 mm	9 mm	8 mm	10 mm	17 mm
OX 1	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm
C 30	13 mm	15 mm	16 mm	9 mm	10 mm	22 mm	24 mm	Resistant	20 mm
F 300	20 mm	16 mm	14 mm	19 mm	13 mm	17 mm	20 mm	21 mm	18 mm

Table 11: Antibiotic susceptibility test results for *S. aureus* collected from human sewage sample

Antibiotics	<i>S. aureus</i> From Human sewage sample							
	<i>S. aureus</i> H1	<i>S. aureus</i> H2	<i>S. aureus</i> H3	<i>S. aureus</i> H4	<i>S. aureus</i> H5	<i>S. aureus</i> H6	<i>S. aureus</i> H7	<i>S. aureus</i> H8
SXT 25	18 mm	22 mm	12 mm	18 mm	17 mm	14 mm	18 mm	21 mm
FOX 30	17 mm	19 mm	14 mm	26 mm	21 mm	18 mm	28 mm	33 mm
PEF 5	24 mm	26 mm	8 mm	13 mm	17 mm	11 mm	15 mm	29 mm
CIP 5	11 mm	16 mm	6 mm	18 mm	21 mm	13 mm	13 mm	25 mm
E15	16 mm	11 mm	7 mm	22 mm	Resistant	8 mm	11 mm	11 mm
CN 10	21 mm	13 mm	5 mm	17 mm	Resistant	6 mm	13 mm	Resistant
K 30	19 mm	11 mm	11 mm	13 mm	19 mm	22 mm	12 mm	20 mm
S 10	24 mm	23 mm	11 mm	16 mm	22 mm	15 mm	28 mm	31 mm
CXM 30	20 mm	21 mm	15 mm	19 mm	26 mm	11 mm	24 mm	Resistant
NA 30	26 mm	21 mm	Resistant	13 mm	Resistant	9 mm	19 mm	17 mm
OX 1	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm
C 30	19 mm	13 mm	28 mm	25 mm	11 mm	7 mm	18 mm	18 mm
F 300	11 mm	9 mm	23 mm	13 mm	Resistant	5 mm	17 mm	22 mm

Table 12: Antibiotic susceptibility test results for *S. aureus* collected from chicken feces

Antibiotics	<i>S. aureus</i> From Chicken Feces							
	<i>S. aureus</i> C1	<i>S. aureus</i> C2	<i>S. aureus</i> C3	<i>S. aureus</i> C4	<i>S. aureus</i> C5	<i>S. aureus</i> C6	<i>S. aureus</i> C7	<i>S. aureus</i> C8
SXT 25	9 mm	11 mm	Resistant	17 mm	21 mm	19 mm	15 mm	22 mm
FOX 30	13 mm	17 mm	12 mm	16 mm	11 mm	25 mm	19 mm	11 mm
PEF 5	11 mm	9 mm	Resistant	28 mm	Resistant	17 mm	15 mm	26 mm
CIP 5	14 mm	11 mm	Resistant	19 mm	Resistant	15 mm	13 mm	15 mm
E15	13 mm	15 mm	Resistant	22 mm	12 mm	18 mm	11 mm	17 mm
CN 10	12 mm	18 mm	Resistant	10 mm	8 mm	14 mm	17 mm	15 mm
K 30	28 mm	13 mm	Resistant	7 mm	19 mm	21 mm	15 mm	13 mm
S 10	24 mm	16 mm	Resistant	24 mm	22 mm	19 mm	21 mm	11 mm
CXM 30	21 mm	19 mm	21 mm	26 mm	23 mm	21 mm	23 mm	17 mm
NA 30	28 mm	26 mm	Resistant	13 mm	17 mm	15 mm	19 mm	24 mm
OX 1	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm	0 mm
C 30	19 mm	16 mm	28 mm	21 mm	31 mm	33 mm	22 mm	25 mm
F 300	14 mm	22 mm	21 mm	11 mm	16 mm	19 mm	17 mm	19 mm




Figure: 8a




Figure: 8b

Figure 8: (a) and (b) Outcome of given antibiotics on *E. coli* B3 isolate collected from bovine sample




Figure: 9a




Figure: 9b

Figure 9: (a) and (b) Outcome of given antibiotics on *E. coli* S3 isolate collected from salad sample




Figure: 10a




Figure: 10b

Figure 10: (a) and (b) Effectiveness of applied antibiotics on *S. aureus* C3 isolate collected from chicken feces




Figure: 11a




Figure: 11b

Figure 11: (a) and (b) Effectiveness of applied antibiotics on *S. aureus* H3 isolate collected from human sewage sample

3.4 Comparative analysis of antibiotic susceptibility profiles of clinical and environmental strains

A comparative study was carried out among clinical and environmental samples isolated from salad sample, human sewage sample, chicken feces, and bovine sample. Results indicate strong differences in the numbers of different environmental isolates and the clinical strains in terms of the development of resistance to standard antibiotic disc.

The clinical strains, its susceptibility to these antibiotics were moderately high. *E. coli* isolates collected from salad sample and bovine sample both showed resistance to two-three different antibiotics each. On the other hand, the clinical strain of *E. coli* was considerably vulnerable to those same antibiotics. In case of *Staphylococcus aureus*, its clinical strain was susceptible to all the antibiotics but one, and yet its environmental isolates collected both from human sewage sample and chicken feces showed resistance to as many as four and eight different antibiotics as presented in the figures below.




Figure 12: Comparison among the efficiency of different antibiotics on clinical and environmental samples of *E. coli* spp collected from bovine sample




Figure 13: Comparison among the efficiency of different antibiotics on clinical and environmental samples of *E. coli* spp collected from salad sample




Figure 14: Comparison among the efficiency of different antibiotics on clinical and environmental samples of *Staphylococcus aureus* spp collected from chicken feces




Figure 15: Comparison among the efficiency of different antibiotics on clinical and environmental samples of *Staphylococcus aureus* spp collected from human sewage sample

3.5 Qualitative analysis of SDS PAGE result:

SDS-PAGE is a reliable method for determining the molecular weight (MW) of an unknown protein, since the migration rate of a protein coated with SDS is inversely proportional to the logarithm of its MW. The key to accurate MW determination is selecting separation conditions that produce a linear relationship between log MW and migration within the likely MW range of the unknown protein.

To ensure accurate MW determination, the protein sample was separated on the same gel with a set of MW standards




Figure 16: SDS PAGE result analysis

The organisms of interest were *E.coli* and *S. aureus*. The clinical sensitive strain and the resistant environmental strain of *E.coli* and *S. aureus* were loaded side by side and distinct band differences in the sensitive and resistant strain were noticed. The unknown protein bands were found only in the resistant strain of the organism. The presence of this unknown protein in the environmental resistant strain can be assumed as the reason of its resistivity as it was not found in the sensitive clinical strain of the same organism.

After separation, the relative migration distance (Rf) of the marker proteins and of the unknown protein was determined. Rf is defined as the mobility of a protein divided by the mobility of the ion front. Because the ion front can be difficult to locate, mobilities are normalized to the tracking dye that migrates only slightly behind the ion front:

$$\text{Rf} = (\text{distance to band}) / (\text{distance to dye front})$$

Table 13: Relative migration distance of protein markers and respective log of MW

Molecular weights of Marker Proteins	Migration distance of the protein (mm)	Migration distance of the dye front (mm)	Rf	Log of MW
120	9	90	0.1	2.08
90	17	90	0.189	1.94
50	34	90	0.45	1.74
35	54	90	0.6	1.56
26	67	90	0.744	1.415
20	80	90	0.889	1.3

Table 14: Relative migration distance of unknown proteins

Micro organisms	Migration distance of the unknown protein (mm)	Migration distance of the dye front (mm)	Rf
<i>E. coli</i>	22	90.5	0.243
<i>S. aureus</i>	41	92.5	0.443

Using the values obtained for the marker proteins, a graph of log MW vs. Rf was plotted. The plot was linear for most proteins, provided they are fully denatured and that the gel percentage is appropriate for the MW range of the sample. The standard curve is sigmoid at extreme MW values because at high MW, the sieving effect of the matrix is so large that molecules are unable to penetrate the gel; at low MW, the sieving effect is negligible and proteins migrate almost freely. The MW of the unknown protein band was determined by interpolating the value from this graph.




Figure: 17

Graph: log MW vs. Rf curve for marker proteins

The results were calculated using this equation below, and the inverse log is the molecular weight of the unknown protein.

$$y = \log \text{ of MW of unknown protein}$$

Table 15: Determining molecular weight of unknown protein by interpolating the linear equation of the graph

Micro organism	Rf of unknown protein	Interpolating the Rf value into the equation	y= log of MW of unknown protein	Inverse log of y (MW of unknown protein in kd)
<i>E. coli</i>	0.243	$y = 0 - 1.2616 * 0.243 + 2.238$	1.931	85.31
<i>S. aureus</i>	0.443	$y = 0 - 1.2616 * 0.443 + 2.238$	1.679	47.753

Chapter 4: Discussion

Antibiotic resistance is the ability of bacteria to endure the antimicrobial activity of antibiotics. It is now apparent that antibiotics that are used to alleviate an infection do not always work anymore. Antibiotic resistance is a global issue, and The US Centers for Disease Control and Prevention (CDC) considers antibiotic resistance one of their peak concerns [7].

One of the greatest discoveries of the 20th century is undoubtedly the finding of antibiotics. This fact is evident, but the genuine speculation is the rise of antibiotic resistance in hospitals, communities, and the environment is associated with their use. The surprising yet alarming new genetic capacities of microorganisms have facilitated from man's overuse of antibiotics to exploit every source of their resistance genes and every means of horizontal gene transmission to develop multi-resistant strains [8]. Such has been seen in this study, that the clinical strain of *Staphylococcus aureus* was significantly vulnerable to the antibiotics, yet the same strain isolated from the feces of chicken showed resistance to as many as eight different antibiotics. This is surely something to be concerned about.

Even prior to penicillin was introduced, resistant strains of bacteria had been detected. The selection pressure has been caused by the exercise of millions of antibiotics over the past 75 years. Ever since antibiotics were introduced to mankind, the abundant use of antibiotics has made almost all disease-causing bacteria resistant to the antibiotics that are commonly used to treat them [7].

Our environment contains numerous products that are man-made or triggered by human contamination, a large variety of examples can be petroleum chemicals, chemical solvents, the products and waste of industrial processes, heavy metals, garbage, and so on. Since the early development of the industrial revolution, humankind has dumped ever-growing amounts of organic and inorganic toxins into streams, rivers, seas, oceans, land, and as well as air. Before the discovery of antibiotics, arsenic, mercury, and iodine were used industrially and medicinally, which under some circumstances, are still employed as such. The major bacterial solution to toxic challenges has taken the form of multivalent pumping systems that prevent intracellular accumulation of structurally diverse bactericidal and bacteriostatic substances [9].

Antimicrobial resistance is a global problem. Although antimicrobial agents have played a major role in reducing the threat from communicable diseases, but the widespread use of these synthetic agents has increasingly resulted in the development of microorganisms that are resistant. In recent years, resistance to these agents has been recognized as a major threat to public health. Emergence of multidrug resistance has limited the therapeutic options, so monitoring the resistance pattern has vital importance [10]. Resistance has increasingly become an even bigger problem in recent years due to the drastically slowed pace at which novel antibiotics are being discovered, while antibiotic use is rising rapidly [11]. This study presents the most important aspects of antibiotic resistance development through a longitudinal study, with the conclusion that it is time to take action. To achieve complete reimbursement of therapeutic applications of antibiotics, we need to gather more information on the rise of antibiotic resistance. Creative approaches to the discovery of novel antibiotics and their accelerated and controlled introduction to therapy are mandatory, given the recent situation of antimicrobial resistance [12].

The environmental strains of *Staphylococcus aureus* and *Escherichia coli* are isolated from different potential environmental sources like salad sample, sewage water, chicken feces, bovine sample etc. Target environmental bacterial species were identified and used for the identification of antibiotic resistance pattern in compare to clinical strains obtained from ICDDR,B (International Center for Diarrheal Disease Research, Bangladesh). All strains were confirmed by means of specific biochemical tests and confirmed in selective media and maintained in nutrient ager medium.

E. coli was also isolated from two different sources for this study, bovine and salad sample. Isolates from both sample showed astounding resistance pattern. Among the isolates from bovine sample, *E. coli* B1 was resistant to PEF5, E15, and S10, *E. coli* B3 was resistant to SXT25, PEF5, and NA30. *E. coli* B5 was resistant to SXT25, and K30. *E. coli* B8 was observed to be resistant to four different antibiotics, which are E15, CN10, CXM30, and C30. All the other isolates from bovine sample showed resistance to none of the tested antibiotics (Figure 8). *E. coli* isolates from salad sample also showed great resistance pattern. *E. coli* S1 was resistant to PEF5, CN10, and CXM30, *E. coli* S3 was resistant to SXT25, and K30, and CXM30. *E. coli* S9 showed

resistance to E15, and K 30. However, no other *E. coli* isolates from salad sample was resistant to any of the test antibiotics (Figure 9).

From chicken feces and human sewage sample, *S. aureus* was isolated. Isolates from both sample showed astonishing resistance pattern. Among the isolates from chicken feces, *S. aureus* C3 was resistant to as many as eight different antibiotics; they are SXT25, PEF5, CIP5, E15, CN10, K30, S10, and NA30 and *S. aureus* C5 was observed to be resistant to PEF5, and CIP5. Other *S. aureus* isolates from chicken feces showed resistance to none of the antibiotics (Figure 10). Among the isolates from human sewage sample, *S. aureus* H3 was resistant to NA30. *S. aureus* H5 showed resistance to four different antibiotics, which are E15, CN10, NA30, and F300. *S. aureus* H8 was observed to be resistant to CN10, and CXM30. Other *S. aureus* isolates collected from human sewage sample showed resistance to none of the test antibiotics (Figure 11). Result of this study shows that these organisms have been well exposed to the tested antimicrobials and they have developed mechanisms to avoid them.

Similar studies were done in various parts of the world to know the resistance pattern of different microorganisms. As example, it is well established that methicillin resistant *Staphylococcus aureus* (MRSA) has been a major challenge for the last 30 years, and in 1993, a study had examined *S. aureus* isolates from 9 different hospitals and found that 15% of isolates were resistant to methicillin [13]. Whereas, in this study *S. aureus* isolated from environmental sample (chicken feces) were found to be resistant to as many as eight different antibiotics, which are SXT25, PEF5, CIP5, E15, CN10, K30, S10, and NA30. This is a drastic change brought on by evolution.

Genetic variability is essential for survival and antimicrobial agents will favor those organisms capable of resisting them. Microorganisms are either inherently resistant, that is, resistance determined by the basic nature of the organism or they can develop acquired resistance, which is the resistance that develops in a previously sensitive strain. Acquired resistance has generally been found to be derived from the use of antimicrobials. In addition, resistance problems are greatest in countries with highest use and in areas where use is concentrated, like intensive therapy units [9].

Antimicrobial resistance pattern monitoring will help us to review the current status of antimicrobial resistance locally, nationally and globally and helpful in minimizing the

consequence of drug resistance, limit the emergence and spread of drug resistant pathogens. This has been a major endeavor of this study. Resistance to antibiotic is increasing and significant community health problems are at risk. An accelerated start up of developing new antibiotics and taking measures to conserve the existing microbial agents can be our way to alleviate the current problem. Also the widespread usage of antibiotics should be brought to a controlled manner along with the measures to help control the bacterial spread to slow the emergence and spread of resistant organisms.

Interpolating the value from (Figure 17), the graph it will give the molecular weight of the unknown protein band. It is also to be noted that the accuracy of this methods in determining the molecular weight of an unknown protein usually ranges from 5% to 10%. The presence of polypeptides for example, lipoproteins and glycol generally leads to erroneous results as they are not fully coated with SDS and therefore, would not behave as expected.

Chapter 5: Conclusion

Undoubtedly, we are living in the wonder years of modern medical sciences and enjoying a longer and healthier life. Antibiotics are one of the major reasons behind this, as it has been our vital source in fighting against various infectious diseases. Since the very beginning of antibiotics starting with the discovery of Penicillin, a diverse range of antibiotics have aided us persistently in our battle against the diseases.

It is alarming that despite all these advances in medical science and development of new antibiotics, the rate of morbidity and mortality due to antibiotic resistant pathogens are increasing. One of many reasons behind this is the widespread antibiotic usage, as it drives the pathogens to develop resistance. Now it is crucial to access the resistance pattern of pathogenic organisms to come up with a solution. This present study has been done in reflection to this fact, and the conclusion can be drawn that this comparative analysis will take us one step closer to resolving this global problem.

References

1. Woolhouse, M.E.J. 2002. Population biology of emerging and re-emerging pathogens. *Trends in Microbiology*. **10** (10).
2. Bonjar, G.H.S., Nik .A.K. and Aghighi. S. 2004. Antibacterial and antifungal survey in plants used indigenous herbal-medicine of south east regions of Iran. *Journal of Biological Sciences*. **4** (3). pp. 405-412.
3. Abiramasundari, P., V. Priya, G.P .Jeyanthi. and Devi. S Gayathri. 2011. Evaluation of the Antibacterial activity of *Cocculus hirsutus*. *Hygeia. J. for Drugs and Med.* **3** (2): pp. 26-31.
4. Lambert, M. 2011. Molecular Biosafety - An update on alternatives to antibiotics - old and new strategies. *Applied Biosafety: ABSA*. **16** (3).
5. Hema, T. A., Arya. A.S, Subha. S, John. R. K and Divya. P.V. 2013. Antimicrobial activity of five south Indian medicinal plants against clinical pathogens. *Int .J. Pharm. Bio. Sci.* **4** (1). pp. 70 – 80.
6. Cappuccino, J.G. and Sherman N. 1996. Microbiology - A Laboratory Manual, 4th Ed. The Benjamin/ Cummings Publishing Co., Inc., Menlo Park, California. pp. 13-182.
7. Antibiotic resistance. 2013. The need for global solutions, LIDC.
8. Fischbach, M.A., and Walsh, C.T. 2010. Antibiotics For Emerging Pathogens. *Science*. **325**. pp. 1089-1093.
9. Davies, J., and Davies, D. 2010. Origins and Evolution of Antibiotic Resistance. *Microbiol Mol Biol Rev.* **74** (3). pp. 417–433.
10. Abdullah, S., J. Gobilik and K. P. Chong. 2013. In Vitro Antimicrobial Activity of *Cynodon dactylon* (L.) Pers. (bermuda) against Selected Pathogens. *International Journal of Pharmacy and Pharmaceutical Sciences*. **4** (5). pp. 227-230.
11. Review on Antimicrobial Resistance. 2014. Antimicrobial Resistance: Tackling a Crisis for the Health and Wealth of Nations.
12. Smith, R.D., and Coast, J. 2002. Antimicrobial resistance: a global response. *Bulletin of the World Health Organization*. **80**. pp. 126-133.
13. Dála, M.M.T. 2001. A Strategy for Control of Antimicrobial Resistance in Ireland. Health and Children.

14. Nathan, C., and Cars, O. 2014. Antibiotic Resistance - Problems, Progress, and Prospects. *The New England Journal of Medicine*. **371**. pp. 1761-1763.
15. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard—Ninth Edition, 2014. CLSI. **32** (2).

APPENDIX-I**Media composition**

The composition of the media used in the present study has been given below. Unless otherwise mentioned, all the media were autoclaved at 121°C for 15 min.

1. Nutrient Agar (Himedia, India)

Ingredients	Amounts (g/L)
Peptic digest of animal tissue	5.0
Beef extract	1.50
Sodium chloride	5.0
Yeast extract	1.50
Agar	15.0

2. Nutrient Broth (Oxoid, England)

Ingredients	Amount (g/L)
Lab-lemco powder	1.0
Yeast extract	2.0
Peptone	5.0
Sodium chloride	5.0

3. Cetrinide agar (Merck, India)

Ingredients	Amount (g/L)
Pancreatic digest of gelatin	20.0
Magnesium chloride hexahydrate	1.4
Potassium sulfate anhydrous	10.0
Cetrinide	0.3
Agar-Agar	13.0

4. T₁N₁ soft agar

Ingredients	Amount (g/L)
Tryptone	0.6 g
Sodium chloride	0.3g
Agar	0.42 g

5. Tryptone soy broth, (Oxoid, England)

Ingredients	Amount (g/L)
Pancreatic digest of Casein	17.0
Papaic digest of soybean meal	3.0
Sodium chloride	5.0
Di-basic potassium phosphate	2.5
Glucose	2.5

6. MacConkey agar (Oxoid, England)

Ingredients	Amount (g/L)
Peptone	20.0
Lactose	10.0
Bile salts	5.0
Sodium chloride	5.0
Neutral red	0.075
Agar	12.0

7. Simmon's citrate agar (Oxoid, England)

Ingredients	Amount (g/L)
Magnesium sulfate	0.2
Ammonium dihydrogen phosphate	0.2
Ammonium phosphate	0.8
Sodium citrate	2.0
Sodium chloride	5.0
Agar	15.0
Bacto brom thymol blue	0.08

8. Peptone Water

Ingredients	Amount (g/L)
Peptone	10.0
Sodium chloride	5.0

9. MR-VP broth

Ingredients	Amount (g/L)
Peptone	7 g
Dextrose	5 g
Potassium phosphate	5 g

10. Triple sugar iron agar (Himedia, India)

Ingredients	Amount (g/L)
Peptic digest of animal tissue	10.0
Sodium chloride	5.0
Lactose	10.0
Sucrose	10.0
Dextrose	1.0
Ferrous sulfate	0.20
Sodium thiosulfate	0.30
Casein enzymatic hydrolysate	10.0
Yeast extract	3.0
Beef extract	3.0

11. Eosine methylene blue agar (Oxoid, England)

Ingredients	Amount (g/L)
Peptone	10.0
Sucrose	5.0
Lactose	5.0
Di-potassium phosphate	2.0
Eosin Y	0.14
Methylene blue	0.065
Agar	13.50

12. Mannitol Salt agar (Oxoid, England)

Ingredients	Amount (g/L)
Peptone	10.0
Manitol	10.0
Lab-lemco powder	1.0
Sodium chloride	75.0
Phenol red	0.025
Agar	15.0

13. Thiosulfate Citrate Bile Salts Sucrose agar (Difco, USA)

Ingredients	Amount (g/L)
Proteose peptone	10.0
Sodium thiosulfate	10.0
Sodium citrate	10.0
Yeast extract	5.0
Oxgall	8.0
Sucrose	20.0
Sodium chloride	10.0
Ferric citrate	1.0
Bromothymol blue	0.04
Thymol blue	0.04
Agar	15.0

14. Xylose Lysine Deoxycholate agar (Himedia, India)

Ingredients	Amount (g/L)
L- lysine	5.0
Lactose	7.50
Sucrose	7.50
Xylose	3.50
Sodium chloride	5.0
Sodium deoxycholate	2.50
Yeast extract	3.0

15. Phenol red (Lactose, Dextrose, Sucrose) Broth

Ingredients	Amount (g/L)
Trypticase	0.4
Lactose	0.2
Sucrose	0.2
Dextrose	0.2
Sodium chloride	0.2
Phenol red	0.00072
Final Ph	7.3

16. Mueller-Hinton agar

Ingredients	Amount (g/L)
Beef, dehydrated infusion from	300.0
Casein hydrolysate	17.5
Starch	1.5
Agar	17.0

APPENDIX-II

Buffers and reagents

- **Phosphate buffered saline (PBS)**

PBS was prepared by dissolving 8.0 g of NaCl, 0.2 g of KCl, 1.44 g of Na₂HPO₄ and 2.0 g of KH₂PO₄ in 800 ml of distilled water. The pH was adjusted to 7.4 with HCl. The final volume was adjusted to 1 liter by distilled water. The solution was sterilized by autoclaving and was stored at room temperature.

- **Kovac's reagent**

5 g of para-dimethylaminobenzaldehyde was dissolved in 75 ml of amyl alcohol. Then concentrated HCl was added to make the final volume 25 ml. This reagent was covered with aluminum foil and stored at 4°C.

- **Methyl red reagent**

0.1 g of methyl red was dissolved in 300 ml of 95% ethyl alcohol. Then distilled water was added to make the final volume 500 ml. This reagent was covered with aluminum foil and stored at 4°C.

- **Barritt's reagent**

Solution A

5 g of alpha-naphthol was dissolved in 95% ethanol. This solution was covered with aluminum foil and stored at 4°C.

Solution B

40 g of KOH was dissolved in distilled water. The solution became warm. After cooling to room temperature, creatine was dissolved by stirring. Distilled water was added. This solution was covered with aluminum foil and stored at

- **Oxidase reagent**

100 mg of N,N,N¹,N¹-tetramethyl-p-phenyldiamine-dihydrochloride was dissolved in 10 ml of distilled water and covered with aluminum foil. Then the solution was stored at 4°C.

- **30% Acrylamide : (for 100ml)**

29.0 g of Acrylamide 1.0gm of Bis- Acrylamide were dissolved in 100ml DH₂O and filtration was needed in every step and using an aluminium foil it was wrapped and stored at 4°C.

- **M Tris (pH 8.8) : (for 100ml)**

18.17 g of 1.5 M Trisma base and 0.4 gm of SDS were dissolved in 100 ml DH₂O rotation was needed O/N. The pH was adjusted then water was added upto level. It was then stored at 4°C

- **0.5 M Tris (pH 6.8) : (for 100ml)**

6.1 g of 1.5 M Trisma base and 0.4 gm SDS was dissolved in 100ml DH₂O and rotation was needed then the pH was adjusted and water was added upto level. Finally stored at 4°C

- **10% SDS Solution: (for 100ml)**

10 g of SDS was dissolved in 100 ml DH₂O it was kept aside as shaking is not suggested. It was then stored at room temperature.

- **10% APS (Ammonium persulphate) : (for 10ml)**

1 g of APS was diluted in 10 ml DH₂O. It was freshly prepared and stored at **-20° C**

- **1x TGS Running Buffer: (for 2.5 L) pH 8.3**

7.5 gm of Trisma base and 36 gm of Glycine were mixed in 25 ml of 10% SDS upto 2.5L of DH₂O.. The pH was adjusted then water is added upto level and finally stored at **4° C**

- **Staining: (for 100 ml)**

10 ml of Acetic Acid and 10 ml of Methanol was added to 80 ml of DH₂O. 0.1-0.2 gm of Comassie Blue was add to it was then filtered and stored at **Room Temperature**

- **Destaining: (for 100 ml)**

10 ml of Acetic Acid and 10 ml of Methanol was diluted in 80 ml of DH₂O. It was then stored at **Room Temperature**

APPENDIX-III

Instruments

The important equipments used through the study are listed below:

❖ Autoclave	SAARC
❖ Freeze (-20°C)	Siemens
❖ Incubator	SAARC
❖ Micropipette (10-100µl)	Eppendorf, Germany
❖ Micropipette (20-200µl)	Eppendorf, Germany
❖ Oven, Model:MH6548SR	LG, China
❖ pH meter, Model: E-201-C	Shanghai Ruosuaa Technology company, China
❖ Refrigerator (4°C), Model: 0636	Samsung
❖ Safety cabinet Class II Microbiological	SAARC
❖ Shaking Incubator, Model: WIS-20R	Daihan Scientific, Korea
❖ Vortex Mixture	VWR International
❖ Water bath	Korea

❖ Weighing balance

ADAM

EQUIPMENT™,

United Kingdom

❖ SDS PAGE Equipment

SERVA

SDS PAGE

Germany